

Recursion Relations for Clebsch-Gordan Coefficients of $U_q(su_2)$ and $U_q(su_{1,1})$

M. Kibler and C. Campigotto

Institut de Physique Nucléaire de Lyon

IN2P3-CNRS et Université Claude Bernard

43 Bd du 11 Novembre 1918, F-69622 Villeurbanne Cedex, France

and

Yu.F. Smirnov¹

Instituto de Física

Universidad Nacional Autónoma de México

México D.F., México

Abstract

We report in this article three- and four-term recursion relations for Clebsch-Gordan coefficients of the quantum algebras $U_q(su_2)$ and $U_q(su_{1,1})$. These relations were obtained by exploiting the complementarity of three quantum algebras in a q -deformation of $sp(8, \mathbb{R})$.

This article constitutes a contribution to the proceedings of the International workshop “Symmetry Methods in Physics” organized by the *Joint Institute for Nuclear Research* (Dubna, Russia, 6-11 July 1993) in memory of Professor Ya.A. Smorodinskiĭ.

¹ *On leave of absence from the Institute of Nuclear Physics, Moscow State University, 119899 Moscow, Russia.*

Recursion Relations for Clebsch-Gordan Coefficients of $U_q(su_2)$ and $U_q(su_{1,1})$

M. Kibler and C. Campigotto

*Institut de Physique Nucléaire de Lyon
IN2P3-CNRS et Université Claude Bernard
43 Bd du 11 Novembre 1918, F-69622 Villeurbanne Cedex, France*

and

Yu.F. Smirnov¹

*Instituto de Física
Universidad Nacional Autónoma de México
México D.F., México*

Abstract

We report in this article three- and four-term recursion relations for Clebsch-Gordan coefficients of the quantum algebras $U_q(su_2)$ and $U_q(su_{1,1})$. These relations were obtained by exploiting the complementarity of three quantum algebras in a q -deformation of $sp(8, \mathbb{R})$.

1. Introduction

The theory of quantum algebras has been the object of numerous investigations both in physics and mathematics. In particular, the one-parameter quantum algebras $U_q(su_2)$ and $U_q(su_{1,1})$ have been investigated by many authors (see, for instance, Refs. [1,2]). In addition, two-parameter deformations of su_2 and u_2 have been worked out in various papers [3-8].

The application to physics of the quantum algebras $U_q(su_2)$ and $U_q(su_{1,1})$ requires the knowledge of the corresponding coupling and recoupling coefficients. Clebsch-Gordan coefficients for $U_q(su_2)$ (and $U_q(su_{1,1})$), in one- and two-parameter formulations, have been calculated by several people (e.g., see Refs. [5,7]). In addition, recursion relations for Clebsch-

¹ *On leave of absence from the Institute of Nuclear Physics, Moscow State University, 119899 Moscow, Russia.*

Gordan coefficients of $U_q(su_2)$ and $U_q(su_{1,1})$ have also been derived, as an extension of the relations corresponding to the $q = 1$ case [9-13], in Refs. [14-16].

It is the aim of this contribution to list three- and four-term recursion relations for the Clebsch-Gordan coefficients of $U_q(su_2)$ derived from an algorithm recently described in Ref. [17]. We shall also give a few three-term recursion relations for the Clebsch-Gordan coefficients of $U_q(su_{1,1})$ derived by means of this algorithm.

This work is dedicated to the memory of the late Professor Ya.A. Smorodinskiĭ who contributed, among many other fields, to the Wigner-Racah algebra of the group SU_2 .

2. The algorithm

The algorithm applied in this article follows from the work of Schwinger [9] and was implicitly used by Rasmussen [11] and Kibler and Grenet [12] in order to derive recursion relations in the $q = 1$ case. In the $q \neq 1$ case, this algorithm has been described in detail by Smirnov and Kibler [17]. We shall not repeat the description of the algorithm in the present paper. However, we shall briefly mention here its main features.

We start from a q -deformation of the dynamical invariance algebra $sp(8, \mathbb{R})$ of the four-dimensional harmonic oscillator. Following the work of Moshinsky and Quesne [18] on the complementarity of Lie groups, we can extract three complementary algebras from two chains of quantum algebras having the q -deformation of $sp(8, \mathbb{R})$ as the head algebra. These algebras are denoted in Ref. [17] as $U_q(su_2^{\mathcal{J}})$, $U_q(su_2^{\Lambda})$ and $U_q(su_{1,1}^{\mathcal{K}})$ and their generators are collectively indicated by $\mathcal{J}$, Λ and $\mathcal{K}$, respectively. The latter generators are built from the four pairs of q -boson operators and the four number operators corresponding to the four-dimensional q -deformed harmonic oscillator. They are defined in such a way to satisfy the co-product rules for the Hopf algebras $U_q(su_2^{\mathcal{J}})$, $U_q(su_2^{\Lambda})$ and $U_q(su_{1,1}^{\mathcal{K}})$ (see Ref. [17]).

The algorithm amounts to calculating, in two different ways, matrix elements of the type $\langle n_1 n_2 n_3 n_4 | X | j : \mu m \kappa \rangle$, where X is either a linear form or a bilinear form of the generators $\{\mathcal{J}\}$, $\{\Lambda\}$ and $\{\mathcal{K}\}$. Furthermore, the vectors $|n_1 n_2 n_3 n_4\rangle$ are state vectors for the four-dimensional harmonic oscillator and the vectors $|j : \mu m \kappa\rangle$ are common eigenstates of $\mathcal{J}^2$, $\mathcal{J}_3$, Λ_3 and $\mathcal{K}_3$, where $\mathcal{J}^2$ stands for the common Casimir operator of $U_q(su_2^{\mathcal{J}})$, $U_q(su_2^{\Lambda})$ and $U_q(su_{1,1}^{\mathcal{K}})$.

It should be emphasized that the algorithm just described also furnishes an elegant way for deriving the q -analogue of Regge symmetries for $U_q(su_2)$ as well as some connecting formulas between Clebsch-Gordan coefficients of $U_q(su_2)$ and $U_q(su_{1,1})$ [17].

3. Recursion relations

We give below recursion relations for the Clebsch-Gordan coefficients of $U_q(su_2)$ and

$U_q(su_{1,1})$ obtained according to the algorithm described in Section 2.

In what follows, we employ the abbreviations

$$\begin{aligned} [x] &= \frac{q^x - q^{-x}}{q - q^{-1}} \\ h_{\pm} &= \frac{1}{2} \pm \frac{1}{2} \quad h_{\mp} = \frac{1}{2} \mp \frac{1}{2} \\ u_{\pm} &= 1 \pm 1 \quad u_{\mp} = 1 \mp 1 \end{aligned}$$

In addition, the Clebsch-Gordan coefficients for $U_q(su_{1,1})$ (for the positive discrete series) and $U_q(su_2)$ are written as $(k_1 k_2 \kappa_1 \kappa_2 | j \kappa)_q$ and $(j_1 j_2 m_1 m_2 | j m)_q$, respectively.

3.1. Recursion relations for $U_q(su_2)$

1. The action of $\mathcal{J}_{\pm}$ on $|j_1 \mp \frac{1}{2}, j_2 \pm \frac{1}{2}, j, m\rangle_q$ leads to

$$\begin{aligned} & \sqrt{[j \mp j_1 \pm j_2 + 1] [j \pm j_1 \mp j_2]} (j_1 j_2 m_1 m_2 | j m)_q \\ &= q^{+\frac{1}{2}(j_1 - m_1 - j_2 + m_2)} \sqrt{[j_1 + m_1 + h_{\mp}] [j_2 + m_2 + h_{\pm}]} (j_1 \mp \frac{1}{2}, j_2 \pm \frac{1}{2}, m_1 \mp \frac{1}{2}, m_2 \pm \frac{1}{2} | j m)_q \\ &+ q^{-\frac{1}{2}(j_1 + m_1 - j_2 - m_2)} \sqrt{[j_1 - m_1 + h_{\mp}] [j_2 - m_2 + h_{\pm}]} (j_1 \mp \frac{1}{2}, j_2 \pm \frac{1}{2}, m_1 \pm \frac{1}{2}, m_2 \mp \frac{1}{2} | j m)_q \end{aligned}$$

2. The action of $\mathcal{K}_{\pm}$ on $|j_1 \mp \frac{1}{2}, j_2 \mp \frac{1}{2}, j, m\rangle_q$ leads to

$$\begin{aligned} & \sqrt{[j_1 + j_2 - j + h_{\mp}] [j_1 + j_2 + j + 1 + h_{\mp}]} (j_1 j_2 m_1 m_2 | j m)_q \\ &= q^{+\frac{1}{2}(j_1 - m_1 + j_2 + m_2 + 1)} \sqrt{[j_1 + m_1 + h_{\mp}] [j_2 - m_2 + h_{\mp}]} (j_1 \mp \frac{1}{2}, j_2 \mp \frac{1}{2}, m_1 \mp \frac{1}{2}, m_2 \pm \frac{1}{2} | j m)_q \\ &- q^{-\frac{1}{2}(j_1 + m_1 + j_2 - m_2 + 1)} \sqrt{[j_1 - m_1 + h_{\mp}] [j_2 + m_2 + h_{\mp}]} (j_1 \mp \frac{1}{2}, j_2 \mp \frac{1}{2}, m_1 \pm \frac{1}{2}, m_2 \mp \frac{1}{2} | j m)_q \end{aligned}$$

3. The action of $\Lambda_{\pm}$ on $|j_1, j_2, j, m \mp 1\rangle_q$ leads to

$$\begin{aligned} & \sqrt{[j \pm m] [j \mp m + 1]} (j_1 j_2 m_1 m_2 | j m)_q \\ &= q^{+m_2} \sqrt{[j_1 \mp m_1 + 1] [j_1 \pm m_1]} (j_1, j_2, m_1 \mp 1, m_2 | j m \mp 1)_q \\ &+ q^{-m_1} \sqrt{[j_2 \mp m_2 + 1] [j_2 \pm m_2]} (j_1, j_2, m_1, m_2 \mp 1 | j m \mp 1)_q \end{aligned}$$

In the limiting case where $q = 1$, the recursion relations 1 to 3 give back well-known relations for the Lie algebra su_2 (see Refs. [12,13]). The six preceding recursion relations are in accordance with the results by Nomura [14], Groza *et al.* [15], Kachurik and Klimyk [15], and Aizawa [16] who derived recursion relations for the Clebsch-Gordan coefficients of $U_q(su_2)$ from q -deformed hypergeometric functions.

We now give twenty four-term recursion relations that can be obtained from twenty operators of type $\mathcal{KK}$, $\mathcal{JJ}$, $\mathcal{JK}$, $\Lambda\Lambda$, $\Lambda\mathcal{K}$ and $\Lambda\mathcal{J}$.

4. The action of $\mathcal{K}_\pm \mathcal{K}_\pm$ on $|j_1 \mp 1, j_2 \mp 1, j, m\rangle_q$ leads to

$$\begin{aligned}
& \sqrt{[j_1 + j_2 + j + 1 + u_\mp] [j_1 + j_2 + j + u_\mp] [j_1 + j_2 - j + u_\mp] [j_1 + j_2 - j \mp 1]} \\
& (j_1 j_2 m_1 m_2 | jm)_q \\
= & q^{+j_1 - m_1 + j_2 + m_2 + 1} \sqrt{[j_1 + m_1 \mp 1] [j_1 + m_1 + u_\mp] [j_2 - m_2 \mp 1] [j_2 - m_2 + u_\mp]} \\
& (j_1 \mp 1, j_2 \mp 1, m_1 \mp 1, m_2 \pm 1 | jm)_q \\
& - [2] q^{m_2 - m_1} \sqrt{[j_1 + m_1 + h_\mp] [j_1 - m_1 + h_\mp] [j_2 + m_2 + h_\mp] [j_2 - m_2 + h_\mp]} \\
& (j_1 \mp 1, j_2 \mp 1, m_1, m_2 | jm)_q \\
& + q^{-j_1 - m_1 - j_2 + m_2 - 1} \sqrt{[j_1 - m_1 \mp 1] [j_1 - m_1 + u_\mp] [j_2 + m_2 \mp 1] [j_2 + m_2 + u_\mp]} \\
& (j_1 \mp 1, j_2 \mp 1, m_1 \pm 1, m_2 \mp 1 | jm)_q
\end{aligned}$$

5. The action of $\mathcal{J}_\pm \mathcal{J}_\pm$ on $|j_1 \mp 1, j_2 \pm 1, j, m\rangle_q$ leads to

$$\begin{aligned}
& \sqrt{[j \mp j_1 \pm j_2 + 2] [j \pm j_1 \mp j_2 - 1] [j \mp j_1 \pm j_2 + 1] [j \pm j_1 \mp j_2]} (j_1 j_2 m_1 m_2 | jm)_q \\
= & q^{+j_1 - m_1 - j_2 + m_2} \sqrt{[j_1 + m_1 \mp 1] [j_1 + m_1 + u_\mp] [j_2 + m_2 \pm 1] [j_2 + m_2 + u_\pm]} \\
& (j_1 \mp 1, j_2 \pm 1, m_1 \mp 1, m_2 \pm 1 | jm)_q \\
& + [2] q^{m_2 - m_1} \sqrt{[j_1 - m_1 + h_\mp] [j_1 + m_1 + h_\mp] [j_2 - m_2 + h_\pm] [j_2 + m_2 + h_\pm]} \\
& (j_1 \mp 1, j_2 \pm 1, m_1, m_2 | jm)_q \\
& + q^{-j_1 - m_1 + j_2 + m_2} \sqrt{[j_1 - m_1 \mp 1] [j_1 - m_1 + u_\mp] [j_2 - m_2 \pm 1] [j_2 - m_2 + u_\pm]} \\
& (j_1 \mp 1, j_2 \pm 1, m_1 \pm 1, m_2 \mp 1 | jm)_q
\end{aligned}$$

6. The action of $\mathcal{J}_\pm \mathcal{K}_\pm$ on $|j_1 \mp 1, j_2, j, m\rangle_q$ leads to

$$\begin{aligned}
& \sqrt{[j_1 + j_2 \mp j + u_\mp] [j_1 + j_2 \pm j + 1] [j \pm j_1 \mp j_2] [j \mp j_1 \pm j_2 + 1]} (j_1 j_2 m_1 m_2 | jm)_q \\
= & q^{+j_1 - m_1 + m_2 + h_\pm} \sqrt{[j_1 + m_1 + u_\mp] [j_1 + m_1 \mp 1] [j_2 \mp m_2] [j_2 \pm m_2 + 1]} \\
& (j_1 \mp 1, j_2, m_1 \mp 1, m_2 \pm 1 | jm)_q \\
& + q^{m_2 - m_1} \sqrt{[j_1 + m_1 + h_\mp] [j_1 - m_1 + h_\mp]} \\
& \left\{ [j_2 - m_2 + h_\pm] q^{+j_2 + h_\mp} - [j_2 + m_2 + h_\pm] q^{-j_2 - h_\mp} \right\} (j_1 \mp 1, j_2, m_1, m_2 | jm)_q \\
& - q^{-j_1 - m_1 + m_2 - h_\pm} \sqrt{[j_1 - m_1 + u_\mp] [j_1 - m_1 \mp 1] [j_2 \pm m_2] [j_2 \mp m_2 + 1]} \\
& (j_1 \mp 1, j_2, m_1 \pm 1, m_2 \mp 1 | jm)_q
\end{aligned}$$

7. The action of $\mathcal{J}_\pm \mathcal{K}_\mp$ on $|j_1, j_2 \pm 1, j, m\rangle_q$ leads to

$$\begin{aligned}
& \sqrt{[j_1 + j_2 \pm j + u_{\pm}] [j_1 + j_2 \mp j + 1] [j \mp j_1 \pm j_2 + 1] [j \pm j_1 \mp j_2]} (j_1 j_2 m_1 m_2 | j m)_q \\
= & -q^{-j_2 + m_2 - m_1 - h_{\mp}} \sqrt{[j_1 \pm m_1] [j_1 \mp m_1 + 1] [j_2 + m_2 + u_{\pm}] [j_2 + m_2 \pm 1]} \\
& (j_1, j_2 \pm 1, m_1 \mp 1, m_2 \pm 1 | j m)_q \\
& + q^{m_2 - m_1} \sqrt{[j_2 + m_2 + h_{\pm}] [j_2 - m_2 + h_{\pm}]} \\
& \left\{ [j_1 + m_1 + h_{\mp}] q^{+j_1 + h_{\pm}} - [j_1 - m_1 + h_{\mp}] q^{-j_1 - h_{\pm}} \right\} \\
& (j_1, j_2 \pm 1, m_1, m_2 | j m)_q \\
& + q^{+j_2 + m_2 - m_1 + h_{\mp}} \sqrt{[j_1 \mp m_1] [j_1 \pm m_1 + 1] [j_2 - m_2 + u_{\pm}] [j_2 - m_2 \pm 1]} \\
& (j_1, j_2 \pm 1, m_1 \pm 1, m_2 \mp 1 | j m)_q
\end{aligned}$$

8. The action of $\Lambda_{\pm} \Lambda_{\pm}$ on $|j_1, j_2, j, m \mp 2\rangle_q$ leads to

$$\begin{aligned}
& \sqrt{[j \pm m - 1] [j \pm m] [j \mp m + 1] [j \mp m + 2]} (j_1 j_2 m_1 m_2 | j m)_q \\
= & q^{+2m_2} \sqrt{[j_1 \pm m_1 - 1] [j_1 \pm m_1] [j_1 \mp m_1 + 1] [j_1 \mp m_1 + 2]} \\
& (j_1, j_2, m_1 \mp 2, m_2 | j, m \mp 2)_q \\
& + [2] q^{m_2 - m_1} \sqrt{[j_1 \pm m_1] [j_1 \mp m_1 + 1] [j_2 \pm m_2] [j_2 \mp m_2 + 1]} \\
& (j_1, j_2, m_1 \mp 1, m_2 \mp 1 | j, m \mp 2)_q \\
& + q^{-2m_1} \sqrt{[j_2 \pm m_2 - 1] [j_2 \pm m_2] [j_2 \mp m_2 + 1] [j_2 \mp m_2 + 2]} \\
& (j_1, j_2, m_1, m_2 \mp 2 | j, m \mp 2)_q
\end{aligned}$$

9. The action of $\Lambda_+ \mathcal{J}_{\pm}$ on $|j_1 \mp \frac{1}{2}, j_2 \pm \frac{1}{2}, j, m - 1\rangle_q$ leads to

$$\begin{aligned}
& \sqrt{[j - m + 1] [j + m] [j \mp j_1 \pm j_2 + 1] [j \pm j_1 \mp j_2]} (j_1 j_2 m_1 m_2 | j m)_q \\
= & q^{\pm \frac{1}{2} (j_1 \mp m_1 - j_2 \pm 3m_2 \pm 1)} \sqrt{[j_1 \pm m_1 \mp 1] [j_1 - m_1 + 1 + h_{\mp}] [j_1 + m_1] [j_2 \pm m_2 + h_{\pm}]} \\
& (j_1 \mp \frac{1}{2}, j_2 \pm \frac{1}{2}, m_1 - \frac{3}{2}, m_2 + \frac{1}{2} | j, m - 1)_q \\
& + \left\{ [j_1 \mp m_1 + h_{\pm}] q^{\mp \frac{1}{2} (j_1 \pm m_1 - j_2 \mp 3m_2 \mp 1)} + [j_2 \pm m_2 + h_{\mp}] q^{\pm \frac{1}{2} (j_1 \mp 3m_1 - j_2 \pm m_2 \mp 1)} \right\} \\
& \sqrt{[j_1 \pm m_1 + h_{\mp}] [j_2 \mp m_2 + h_{\pm}]} (j_1 \mp \frac{1}{2}, j_2 \pm \frac{1}{2}, m_1 - \frac{1}{2}, m_2 - \frac{1}{2} | j, m - 1)_q \\
& + q^{\mp \frac{1}{2} (j_1 \pm 3m_1 - j_2 \mp m_2 \pm 1)} \sqrt{[j_2 \mp m_2 \pm 1] [j_2 - m_2 + 1 + h_{\pm}] [j_2 + m_2] [j_1 \mp m_1 + h_{\mp}]} \\
& (j_1 \mp \frac{1}{2}, j_2 \pm \frac{1}{2}, m_1 + \frac{1}{2}, m_2 - \frac{3}{2} | j, m - 1)_q
\end{aligned}$$

10. The action of $\Lambda_+ \mathcal{K}_{\pm}$ on $|j_1 \mp \frac{1}{2}, j_2 \mp \frac{1}{2}, j, m - 1\rangle_q$ leads to

$$\begin{aligned}
& \sqrt{[j-m+1][j+m][j_1+j_2\pm j+1][j_1+j_2\mp j+u_{\mp}]}(j_1j_2m_1m_2|jm)_q \\
&= \pm q^{\pm\frac{1}{2}(j_1\mp m_1+j_2\pm 3m_2+u_{\pm})} \sqrt{[j_1+m_1][j_1\pm m_1\mp 1][j_1-m_1+1+h_{\mp}][j_2\mp m_2+h_{\mp}]} \\
&\quad (j_1\mp \frac{1}{2}, j_2\mp \frac{1}{2}, m_1-\frac{3}{2}, m_2+\frac{1}{2}|j, m-1)_q \\
&\quad \pm \left\{ [j_2\mp m_2+h_{\pm}] q^{\pm\frac{1}{2}(j_1\mp 3m_1+j_2\pm m_2+u_{\mp})} - [j_1\mp m_1+h_{\pm}] q^{\mp\frac{1}{2}(j_1\pm m_1+j_2\mp 3m_2+u_{\mp})} \right\} \\
&\quad \sqrt{[j_1\pm m_1+h_{\mp}][j_2\pm m_2+h_{\mp}]}(j_1\mp \frac{1}{2}, j_2\mp \frac{1}{2}, m_1-\frac{1}{2}, m_2-\frac{1}{2}|j, m-1)_q \\
&\quad \mp q^{\mp\frac{1}{2}(j_1\pm 3m_1+j_2\mp m_2+u_{\pm})} \sqrt{[j_2+m_2][j_2\pm m_2\mp 1][j_2-m_2+1+h_{\mp}][j_1\mp m_1+h_{\mp}]} \\
&\quad (j_1\mp \frac{1}{2}, j_2\mp \frac{1}{2}, m_1+\frac{1}{2}, m_2-\frac{3}{2}|j, m-1)_q
\end{aligned}$$

11. The action of $\Lambda_- \mathcal{J}_{\pm}$ on $|j_1\mp \frac{1}{2}, j_2\pm \frac{1}{2}, j, m+1\rangle_q$ leads to

$$\begin{aligned}
& \sqrt{[j-m][j+m+1][j\mp j_1\pm j_2+1][j\pm j_1\mp j_2]}(j_1j_2m_1m_2|jm)_q \\
&= q^{\pm\frac{1}{2}(j_1\mp 3m_1-j_2\pm m_2\pm 1)} \sqrt{[j_2-m_2][j_2\pm m_2\pm 1][j_2+m_2+1+h_{\pm}][j_1\pm m_1+h_{\mp}]} \\
&\quad (j_1\mp \frac{1}{2}, j_2\pm \frac{1}{2}, m_1-\frac{1}{2}, m_2+\frac{3}{2}|j, m+1)_q \\
&\quad + \left\{ [j_1\pm m_1+h_{\pm}] q^{\pm\frac{1}{2}(j_1\mp m_1-j_2\pm 3m_2\mp 1)} + [j_2\mp m_2+h_{\mp}] q^{\mp\frac{1}{2}(j_1\pm 3m_1-j_2\mp m_2\mp 1)} \right\} \\
&\quad \sqrt{[j_1\mp m_1+h_{\mp}][j_2\pm m_2+h_{\pm}]}(j_1\mp \frac{1}{2}, j_2\pm \frac{1}{2}, m_1+\frac{1}{2}, m_2+\frac{1}{2}|j, m+1)_q \\
&\quad + q^{\mp\frac{1}{2}(j_1\pm m_1-j_2\mp 3m_2\pm 1)} \sqrt{[j_1-m_1][j_1\mp m_1\mp 1][j_1+m_1+1+h_{\mp}][j_2\mp m_2+h_{\pm}]} \\
&\quad (j_1\mp \frac{1}{2}, j_2\pm \frac{1}{2}, m_1+\frac{3}{2}, m_2-\frac{1}{2}|j, m+1)_q
\end{aligned}$$

12. The action of $\Lambda_- \mathcal{K}_{\pm}$ on $|j_1\mp \frac{1}{2}, j_2\mp \frac{1}{2}, j, m+1\rangle_q$ leads to

$$\begin{aligned}
& \sqrt{[j-m][j+m+1][j_1+j_2\mp j+u_{\mp}][j_1+j_2\pm j+1]}(j_1j_2m_1m_2|jm)_q \\
&= \pm q^{\pm\frac{1}{2}(j_1\mp 3m_1+j_2\pm m_2+u_{\pm})} \sqrt{[j_1\pm m_1+h_{\mp}][j_2-m_2][j_2\mp m_2\mp 1][j_2+m_2+1+h_{\mp}]} \\
&\quad (j_1\mp \frac{1}{2}, j_2\mp \frac{1}{2}, m_1-\frac{1}{2}, m_2+\frac{3}{2}|j, m+1)_q \\
&\quad \pm \left\{ [j_1\pm m_1+h_{\pm}] q^{\pm\frac{1}{2}(j_1\mp m_1+j_2\pm 3m_2+u_{\mp})} - [j_2\pm m_2+h_{\pm}] q^{\mp\frac{1}{2}(j_1\pm 3m_1+j_2\mp m_2+u_{\mp})} \right\} \\
&\quad \sqrt{[j_1\mp m_1+h_{\mp}][j_2\mp m_2+h_{\mp}]}(j_1\mp \frac{1}{2}, j_2\mp \frac{1}{2}, m_1+\frac{1}{2}, m_2+\frac{1}{2}|j, m+1)_q \\
&\quad \mp q^{\mp\frac{1}{2}(j_1\pm m_1+j_2\mp 3m_2+u_{\pm})} \sqrt{[j_2\pm m_2+h_{\mp}][j_1-m_1][j_1\mp m_1\mp 1][j_1+m_1+1+h_{\mp}]} \\
&\quad (j_1\mp \frac{1}{2}, j_2\mp \frac{1}{2}, m_1+\frac{3}{2}, m_2-\frac{1}{2}|j, m+1)_q
\end{aligned}$$

13. The action of $\Lambda_{\pm} \Lambda_{\mp}$ on $|j_1, j_2, j, m\rangle_q$ leads to

$$\begin{aligned}
& [j\pm m][j\mp m+1](j_1j_2m_1m_2|jm)_q \\
&= q^{m_2-m_1+1} \sqrt{[j_1-m_1+1][j_1+m_1][j_2+m_2+1][j_2-m_2]}(j_1, j_2, m_1-1, m_2+1|jm)_q \\
&\quad + \left\{ q^{+2m_2}[j_1\mp m_1+1][j_1\pm m_1] + q^{-2m_1}[j_2\mp m_2+1][j_2\pm m_2] \right\} (j_1j_2m_1m_2|jm)_q \\
&\quad + q^{m_2-m_1-1} \sqrt{[j_1+m_1+1][j_1-m_1][j_2-m_2+1][j_2+m_2]}(j_1, j_2, m_1+1, m_2-1|jm)_q
\end{aligned}$$

In the limiting case where $q = 1$, the recursion relations 4 to 12 are in agreement with the results of Kibler and Grenet [12].

3.2. Recursion relations for $U_q(su_{1,1})$

14. The action of $\mathcal{J}_\pm$ on $|k_1 \mp \frac{1}{2}, k_2 \pm \frac{1}{2}, j, \kappa\rangle_q$ leads to

$$\begin{aligned} & \sqrt{[j \mp k_1 \pm k_2 + 1] [j \pm k_1 \mp k_2]} (k_1 k_2 \kappa_1 \kappa_2 |j\kappa)_q \\ &= q^{\frac{1}{2}(\pm\kappa_1 \mp \kappa_2 - k_1 + k_2)} \sqrt{[\kappa_1 \pm k_1] [\kappa_2 \pm k_2 \pm 1]} (k_1 - \frac{1}{2}, k_2 + \frac{1}{2}, \kappa_1 \mp \frac{1}{2}, \kappa_2 \pm \frac{1}{2} |j\kappa)_q \\ &+ q^{\frac{1}{2}(\mp\kappa_1 \pm \kappa_2 - k_1 + k_2)} \sqrt{[\kappa_2 \mp k_2] [\kappa_1 \mp k_1 \mp 1]} (k_1 + \frac{1}{2}, k_2 - \frac{1}{2}, \kappa_1 \mp \frac{1}{2}, \kappa_2 \pm \frac{1}{2} |j\kappa)_q \end{aligned}$$

15. The action of $\mathcal{K}_\pm$ on $|k_1, k_2, j, \kappa \mp 1\rangle_q$ leads to

$$\begin{aligned} & \sqrt{[\kappa \mp j \mp 1] [\kappa \pm j]} (k_1 k_2 \kappa_1 \kappa_2 |j\kappa)_q \\ &= \pm q^{\frac{1}{2}(\pm\kappa_1 \pm \kappa_2 - k_1 + k_2)} \sqrt{[\kappa_1 \pm k_1] [\kappa_2 \mp k_2 \mp 1]} (k_1 - \frac{1}{2}, k_2 + \frac{1}{2}, \kappa_1 \mp \frac{1}{2}, \kappa_2 \mp \frac{1}{2} |j, \kappa \mp 1)_q \\ &\mp q^{\frac{1}{2}(\mp\kappa_1 \mp \kappa_2 - k_1 + k_2)} \sqrt{[\kappa_2 \pm k_2] [\kappa_1 \mp k_1 \mp 1]} (k_1 + \frac{1}{2}, k_2 - \frac{1}{2}, \kappa_1 \mp \frac{1}{2}, \kappa_2 \mp \frac{1}{2} |j, \kappa \mp 1)_q \end{aligned}$$

16. The action of $\Lambda_\pm$ on $|k_1 \mp \frac{1}{2}, k_2 \mp \frac{1}{2}, j, \kappa\rangle_q$ leads to

$$\begin{aligned} & \sqrt{[j \pm k_1 \pm k_2 \pm 1] [j \mp k_1 \mp k_2 + u_\mp]} (k_1 k_2 \kappa_1 \kappa_2 |j\kappa)_q \\ &= q^{+\frac{1}{2}(k_1 + k_2 - \kappa_1 + \kappa_2 + u_\mp)} \sqrt{[\kappa_1 - k_1 - h_\mp] [\kappa_1 + k_1 + h_\mp]} (k_1 \mp 1, k_2, \kappa_1, \kappa_2 |j\kappa)_q \\ &+ q^{-\frac{1}{2}(k_1 + k_2 + \kappa_1 - \kappa_2 + u_\mp)} \sqrt{[\kappa_2 - k_2 - h_\mp] [\kappa_2 + k_2 + h_\mp]} (k_1, k_2 \mp 1, \kappa_1, \kappa_2 |j\kappa)_q \end{aligned}$$

The recursion relations 14 to 16 are in accordance with the results by Nomura [14], Groza *et al.* [15], Kachurik and Klimyk [15], and Aizawa [16] who derived recursion relations for the Clebsch-Gordan coefficients of $U_q(su_{1,1})$ from q -deformed hypergeometric functions.

Acknowledgements

One of the authors (Yu.F. S.) is grateful to the *Institut de Physique Nucléaire de Lyon* for the hospitality extended to him during his stay in Lyon-Villeurbanne where this work was completed.

References

- [1] Biedenharn L.C., J. Phys. A: Math. Gen. **22** (1989) L873.
- [2] Macfarlane A.J., J. Phys. A: Math. Gen. **22** (1989) 4581.
- [3] Chakrabarti R. and Jagannathan R., J. Phys. A: Math. Gen. **24** (1991) L711.
- [4] Schirrmacher A., Wess J. and Zumino B., Z. Phys. C **49** (1991) 317.
- [5] Smirnov Yu.F. and Wehrhahn R.F., J. Phys. A: Math. Gen. **25** (1992) 5563.
- [6] Kibler M.R., in *Symmetry and Structural Properties of Condensed Matter*, Eds., Florek W., Lipinski D. and Lulek T., World Scientific, Singapore (1993), p. 445.
- [7] Dorešić M., Meljanac S. and Mileković M., Fizika B **2** (1993) 43.
- [8] Barbier R., Meyer J. and Kibler M., Preprint LYCEN 9348.
- [9] Schwinger J., Report U.S. Atom. Ener. Comm. NYO-3071 (1952), published in *Quantum Theory of Angular Momentum*, Eds., Biedenharn L.C. and van Dam H., Academic Press, New York (1965).
- [10] Smorodinskiĭ Ya.A. and Shelepin L.A., Usp. Fiz. Nauk **106** (1972) 3. (English translation: Usp. Fiz. Nauk. **15** (1972) 1.)
- [11] Rasmussen W., J. Phys. A: Math. Gen. **8** (1975) 1038.
- [12] Kibler M. and Grenet G., J. Math. Phys. **21** (1980) 422.
- [13] Biedenharn L.C. and Louck J.D., in *Encyclopedia of Mathematics and Its Applications*, Ed., Rota G.-C., Addison-Wesley, Reading, Massachusetts (1981), Vol. 8/9.
- [14] Nomura M., J. Phys. Soc. Japan **59** (1990) 1954 ; Nomura M., J. Phys. Soc. Japan **59** (1990) 2345.
- [15] Groza V.A., Kachurik I.I. and Klimyk A.U., J. Math. Phys. **31** (1990) 2769 ; Kachurik I.I. and Klimyk A.U., J. Phys. A: Math. Gen. **24** (1991) 4009.
- [16] Aizawa N., J. Math. Phys. **34** (1993) 1937.
- [17] Smirnov Yu.F. and Kibler M.R., in *Symmetries in Science VI: From the Rotation Group to Quantum Algebras*, Ed., Gruber B., Plenum Press, New York (1993), p. 691.
- [18] Moshinsky M. and Quesne C., Phys. Lett. **B 29** (1969) 482 ; Moshinsky M. and Quesne C., J. Math. Phys. **11** (1970) 1631.